Effector Biology of Plant-Associated Organisms: Concepts and Perspectives

J. WIN,¹ A. CHAPARRO-GARCIA,¹ K. BELHAJ,¹ D.G.O. SAUNDERS,¹ K. YOSHIDA,¹ S. DONG,¹ S. SCHORNACK,¹ C. ZIPFEL,¹ S. ROBATZEK,¹ S.A. HOGENHOUT,² AND S. KAMOUN¹

¹*The Sainsbury Laboratory, Norwich Research Park, Norwich, NR4 7UH United Kingdom*

²*Department of Cell and Developmental Biology, The John Innes Centre, Norwich Research Park, Norwich, NR4 7UH United Kingdom*

Correspondence: sophien.kamoun@tsl.ac.uk

Every plant is closely associated with a variety of living organisms. Therefore, deciphering how plants interact with mutualistic and parasitic organisms is essential for a comprehensive understanding of the biology of plants. The field of plant–biotic interactions has recently coalesced around an integrated model. Major classes of molecular players both from plants and their associated organisms have been revealed. These include cell surface and intracellular immune receptors of plants as well as apoplastic and host-cell-translocated (cytoplasmic) effectors of the invading organism. This article focuses on effectors, molecules secreted by plant-associated organisms that alter plant processes. Effectors have emerged as a central class of molecules in our integrated view of plant–microbe interactions. Their study has significantly contributed to advancing our knowledge of plant hormones, plant development, plant receptors, and epigenetics. Many pathogen effectors are extraordinary examples of biological innovation; they include some of the most remarkable proteins known to function inside plant cells. Here, we review some of the key concepts that have emerged from the study of the effectors of plant-associated organisms. In particular, we focus on how effectors function in plant tissues and discuss future perspectives in the field of effector biology.

Interactions with the biota are critical to plant life. In nature, every single plant is closely associated with a variety of living organisms, particularly microbes (Hines and Zahn 2009; Rodriguez et al. 2009; Redford et al. 2010). Plants are repeatedly attacked by a multitude of pathogens and pests, including viruses, bacteria, fungi, oomycetes, nematodes, and insects (Agrios 2005). Some plants can also establish parasitic relationships with other plants (Westwood et al. 2010). Given the ubiquity of plant-associated organisms in nature, deciphering how plants interact with mutualistic and parasitic organisms is essential for a comprehensive understanding of the biology of plants.

Despite the diversity and multitude of organisms that interact with plants, a series of common notions define our understanding of plant–biotic interactions. The field of plant–biotic interactions has recently coalesced around an integrated model that applies to all types of pathogens and mutualists (Dodds and Rathjen 2010). Both plants and associated organisms contribute molecular players that dictate the outcome of the plant–biotic interaction. For example, plants contribute cell surface and intracellular immune receptors, and the colonizing organisms produce a repertoire of effectors that modulate plant processes, including the induction and suppression of plant defenses (Fig. 1). This article focuses on effectors, molecules secreted by plant-associated organisms that alter host-cell structure and function (Hogenhout et al. 2009).

Effectors have generally evolved to enable parasitism, for example, by suppressing plant immunity or by mod-

ifying plant physiology to support growth and spread of the parasite. These responses are collectively known as effector-triggered susceptibility (ETS) (Fig. 1A) and are achieved through the perturbation of a set of host processes that we define as effector-targeted pathways (ETPs) (Table 1). However, effectors can also “trip the wire” and activate plant immune receptors, a response known as effector-triggered immunity (ETI), when it is mediated by intracellular immunoreceptors of the nucleotide-binding leucine-rich repeat (NB-LRR) class (Fig. 1B). In contrast, plant cell surface receptors recognize apoplastic effectors as well as conserved pathogen molecules known as pathogen-associated molecular patterns (PAMPs) (Fig. 1B) (Boller and Felix 2009; Thomma et al. 2011). These receptors, which are typically receptor kinases and receptor-like proteins, are collectively known as pattern recognition receptors (PRRs). The resulting immune response is known as PRR-triggered immunity or PAMP-triggered immunity (PTI).

Many pathogen effectors have evolved to suppress PTI, providing additional twists to the arms race co-evolution between pathogens and plants (Chisholm et al. 2006; Jones and Dangl 2006; Dodds and Rathjen 2010). Effectors are therefore targeted by dynamic and sometimes opposite evolutionary forces because they can have either positive or negative fitness effects on the pathogen or pest, depending on the host plant genotype. As a consequence, effector genes are among the most rapidly evolving genes in the genome of plant-associated organisms.

Figure 1. The concept of effectors in plant immunity. Infectious pathogens such as bacteria, fungi, oomycete, and nematodes deliver effectors at the interface of the host plant (apoplasmic effectors, AE) or inside the cell (cytoplasmic effectors, CE). Host-translocated (cytoplasmic) effectors are delivered into the host cytoplasm through a type-III secretion pilus or specialized infectious structures called haustoria that form within the cell. Pathogen effectors traffic to various compartments, bind, and manipulate different host proteins called targets. Depending on their localization in the cells, these targets are designated as apoplasmic effector target (AET) and cytoplasmic effector target (CET). Effector–target interactions impact the outcome of the interaction between the pathogen and its host. In susceptible genotypes (*A*), these molecular interactions can alter plant cell processes and suppress immune responses, leading to effector-triggered susceptibility (ETS) and host colonization. In resistant genotypes (*B*), these interactions are perceived by key sensing receptors of the immune system that, in turn, stop pathogen growth. Cell surface pattern recognition receptors (PRRs) detect pathogen-associated molecular patterns (PAMPs), apoplasmic effectors, and/or apoplasmic effector–target interactions to initiate PRR-triggered immunity (PTI). Intracellular nucleotide-binding receptors (NB-LRR) induce NB-LRR-triggered immunity (ETI) on recognition of cytoplasmic effectors and/or cytoplasmic effectors–target interactions. (Modified from Dodds and Rathjen 2010.)

Effectors have emerged as a central class of molecules in our integrated view of plant–microbe interactions. Understanding the molecular functions of effectors is widely accepted as essential for a mechanistic understanding of the processes underlying plant colonization by invading organisms. Effectors can also serve as molecular probes for unraveling key plant processes. Their study has significantly contributed to advancing our knowledge of plant hormones, plant development, plant receptors, signal transduction pathways, and epigenetics. Many

pathogen effectors are extraordinary examples of biological innovation; they include some of the most remarkable proteins known to function inside plant cells. Among such remarkable effectors, we highlight the transcription activator-like (TAL) effectors of *Xanthomonas* bacteria, modular proteins that bind specific plant promoter elements to activate plant gene transcription (Boch et al. 2009); the bacterial effectors AvrPtoB and coronatine, which mimic plant E3 ubiquitin ligases (Janjusevic et al. 2006; Rosebrock et al. 2007) and the hormone

Table 1. Definition of terms used in this manuscript

Term	Definition
Apoplasmic effector	Effectors that are secreted to the host extracellular space. They interact with host extracellular proteins and can also be recognized by PRRs.
Cytoplasmic (host-translocated) effector	Effectors that are translocated inside the host cell through specialized microbial structures such as the type-III secretion system or infection vesicles and haustoria. Inside the host cell, they traffic to different subcellular compartments where they exert their functions.
Effector	Microbial and pest secreted molecules that alter host-cell processes or structure generally promoting the microbe lifestyle. Effector functions are as diverse as suppressing immune responses or enhancing access to nutrients.
Effector helper	Host molecules genetically upstream of the effector-mediated target modulation that are required by the effector to exert its function but do not necessarily have a direct impact on the cellular process that the effector is targeting.
Effector target	Host molecules that the effector modifies to manipulate host processes, thus enhancing the microbe fitness and ability to colonize and spread to other hosts in susceptible plants.
Effector-targeted pathway (ETP)	Natural cellular physiological processes such as RNA silencing, innate immunity, intracellular trafficking, cell signaling, and transcription that are manipulated by effectors for the pathogen benefit.
Effector-triggered susceptibility (ETS)	The outcome of the deployment of effector molecules that favors pathogen virulence. ETS can be the result of suppressing innate immunity or enhancing the pathogen nutrient uptake.
Host susceptibility factor	Host molecules that are hijacked by the invading microbe to support effector functions. Both host targets and host helpers are susceptibility factors; they operate in conjunction to the advantage of the pathogen.
Nucleotide-binding leucine-rich repeat (NB-LRR) proteins	Intracellular (cytoplasmic) receptor proteins specific of plants that have evolved to recognize pathogen effectors or their impact on cellular processes. They contain a nucleotide-binding site (NB) and leucine-rich repeat (LRR) domains coupled to either coiled-coil (CC) or Toll/interleukin-1 receptor (TIR) motifs in the amino-terminal region. NB-LRR receptor activation usually leads to a cell death response known as the hypersensitive response (HR).
Pathogen-associated molecular patterns (PAMPs)	Microbial-derived signatures that are conserved across an entire class of microbes and that contribute to organism fitness. These molecules are often present in nonpathogenic microorganisms; therefore, they are also known as microbe-associated molecular patterns (MAMPs)
Pattern recognition receptors (PRRs)	Cell surface localized receptors that recognize conserved microbial patterns (PAMPs) or apoplasmic effectors. PRRs allow the plant to discriminate between self and nonself molecules given their ability to sense microbial molecules. There are two classes of PRRs: receptor-like kinases (RLKs) and receptor-like proteins (RLPs). The extracellular domains that mediate recognition comprise LRRs or LysM motifs.

jasmonoyl-isoleucine (JA-Ile) (Weiler et al. 1994; Bender et al. 1999), respectively; SAP11 of phytoplasma bacteria that binds host TCP transcription factors and destabilizes CIN-TCPs to alter plant development and inhibit jasmonate (JA) synthesis (Sugio et al. 2011); and the viral protein P19 that binds double-stranded RNA with high affinity in order to suppress RNA interference (RNAi) in host plant cells (Lakatos et al. 2004).

This article summarizes and discusses some of the key concepts that have emerged from the study of the effectors of plant-associated organisms. In particular, we focus on how effectors function in plant tissues and discuss future perspectives in the field of effector biology.

EFFECTORS: USAGE AND DEFINITION

The term “effector” has become widely used in the field of plant–biotic interactions, beginning ~12 years ago (for review, see Hogenhout et al. 2009). This rise in usage was in part because the neutral term “effector” addresses the conceptual limitations of teleological terminology such as “avirulence” and “elicitor.” These outdated terms misrepresented the fact that the same

molecules can have both positive and negative effects on the pathogen depending on the host genotype (see discussions in Hogenhout et al. 2009). Plant bacteriologists initially equated effectors to the proteins delivered inside plant cells via the type-III secretion system (T3SS). However, the usage broadened once it became evident that pathogens use a multitude of mechanisms to translocate proteins inside plant cells. More recently, the term effector has also been extended to secreted proteins of mutualistic microbes that contribute to the establishment of a symbiotic relationship with plants (Kloppholz et al. 2011; Plett et al. 2011).

As previously discussed, we favor a broad, inclusive definition of effectors as “molecules secreted by plant-associated organisms that alter host-cell structure and function” (Hogenhout et al. 2009). This broader definition is particularly suitable when only limited information about the function of the “effector” molecules is known, for instance, in the case of effectors with avirulence or elicitor activities, that is, that trigger plant defense responses. Once more information about the functions of an “effector” is revealed, it is preferable to use descriptors that reflect these specific activities (for instance, protease inhibitor).

Because of the prominence of immunity in the study of plant–microbe interactions, there is a perceptible tendency in the literature to assume that effectors always function as immunosuppressors. This view is too narrow; there is more to effector function than suppression of immunity. Although it appears that the majority of *Pseudomonas syringae* T3SS effectors function as immunosuppressors, many other distinct functions have been assigned to pathogen effectors. The TAL effectors of xanthomonads activate host genes involved in processes as diverse as cell size, sugar transport, and epigenetics (Duan et al. 1999; Chen et al. 2010; Domingues et al. 2010; de Souza et al. 2012). Aster yellows phytoplasmas secrete SAP proteins that directly perturb plant development to increase the mass of host green tissue and the likelihood of transmission by their insect vector (MacLean et al. 2011; Sugio et al. 2011). Surely, many more diverse functions of effectors remain to be discovered.

Another issue worth highlighting is the case of necrotrophs, pathogens that colonize dead plant tissues. Many necrotrophs secrete toxin effectors that actively cause tissue necrosis thus contributing to pathogen virulence (Mengiste 2012). Paradoxically, some necrotrophic fungi kill plants by activating immune receptors via secreted toxin effectors. In one well-characterized example, the *Arabidopsis* *LOV1* gene, a member of the NB-LRR resistance gene family, was shown to determine susceptibility to the fungus *Cochliobolus victoriae* and sensitivity to the toxin victorin (Lorang et al. 2007). In this case, the tables are turned—the immunoreceptor becomes a liability for the plant, contributing to susceptibility to a fungal disease.

EFFECTORS TRAFFIC TO DIFFERENT CELLULAR COMPARTMENTS IN PLANT CELLS

Microbes and other plant-associated organisms secrete effectors to different sites in the host plant. Several pathogens, including fungi and oomycetes, colonize plant intercellular spaces and secrete effectors that target defenses in the plant apoplast (Fig. 1A). Classic examples include the effectors of the dothidiomycete leaf mold fungus *Cladosporium fulvum* Avr2, Avr4, and ECP6, which target various extracellular processes of the host plant tomato (van den Burg et al. 2006; Bolton et al. 2008; van Esse et al. 2008). Whereas Avr2 is an inhibitor of tomato apoplastic cysteine proteases, ECP6 interferes with the perception of *C. fulvum* cell wall chitin by tomato cell surface immune receptors (de Jonge et al. 2010). Other microbes deliver their effectors inside plant cells, typically through specialized structures (Fig. 1A). Several fungi and oomycetes use hyphal extensions called haustoria to push inside plant cells, where they remain enveloped with a plant-derived membrane, resulting in a close interface through which effectors are thought to traffic inside host cells (Dodds et al. 2009; de Jonge et al. 2011). Effectors of the rice blast fungus

Magnaporthe oryzae accumulate and translocate into host cells at a membranous cap known as the biotrophic interfacial complex (BIC) (Valent and Khang 2010). Bacteria rely on specialized secretion pili, such as pseudomonad and xanthomonad T3SSs (Buttner and He 2009). The crown gall bacterial pathogen *Agrobacterium tumefaciens* and several related species use a type-IV secretion apparatus to transfer a DNA fragment (T-DNA) inside plant cells in complex with bacterial virulence proteins such as VirD2 and VirE2 (Christie 2004). The T-DNA stably inserts into the plant genome to express biosynthetic enzymes for plant hormones and amino acids, resulting in the formation of tumors and production of bacterial nutrients. Phytoplasmas, plant pathogenic bacteria that, similarly to animal parasitic mycoplasmas diverged from Gram-positive bacteria through loss of outer cell wall and dramatic genome reductions, are intracellular obligate pathogens of plants (Hogenhout and Loria 2008). They directly secrete effector proteins inside host plant cytoplasm using the general Sec-dependent (type II) secretory pathway, and, unlike extracellular bacterial phytopathogens, do not appear to require specialized secretory systems for pathogenesis (Kakizawa et al. 2004).

Once translocated into the plant cytoplasm, effectors can traffic to different subcellular compartments, including organelles and various membrane compartments. A large number of effectors accumulate in the plant nucleus (see also Caillaud et al. 2012, this volume). Cytoplasmic effectors, such as xanthomonad TAL effectors, phytoplasma SAP11, and *Phytophthora* Crinklers (CRNs), carry nuclear localization signals (NLSs) enabling them to subvert the plant protein importin- α to mediate import into the plant nucleus (Van den Ackerveken et al. 1996; Bai et al. 2009; Schornack et al. 2010). The *P. syringae* type-III effector HopG1 targets plant mitochondria (Block et al. 2010), whereas HopI1 localizes to plant chloroplasts, where it suppresses salicylic acid synthesis and disrupts the thylakoid stack structure (Jelenska et al. 2007). Another *P. syringae* type-III effector, the acetyltransferase HopZ1a, targets the plant cytoskeleton (Lee et al. 2012a). HopZ1a disrupts microtubule networks, subsequently interfering with the plant secretory pathway and suppressing cell-wall-mediated defense (Lee et al. 2012a). Recently, some oomycete effectors, notably *Phytophthora infestans* AVRblb2 and AVR2, were shown to relocate from the plasma membrane to accumulate around haustoria inside infected plant cells (Bozkurt et al. 2011; Saunders et al. 2012). This suggests that these effectors associate with dynamic focal processes that take place at pathogen penetration sites (Underwood and Somerville 2008). T3SS effectors of the plant pathogenic bacteria *P. syringae* and *Xanthomonas campestris* have been reported to target secretory pathways, suggesting that interference with polarized vesicle trafficking is a common pathogenesis process (Bartetzko et al. 2009; Kaffarnik et al. 2009; Nomura et al. 2011). One example is the *P. syringae* effector HopM1 that localizes to the *trans*-Golgi network/early endosome to destabilize the host protein AtMIN7, also known as BEN1 regulating

endosomal recycling (Tanaka et al. 2009; Nomura et al. 2011).

TEMPORAL ASPECTS OF PATHOGEN DEPLOYMENT OF EFFECTORS

Effectors not only need to be at the right location during infection but they also require deployment at the right time for successful colonization. Indeed, pathogens secrete different sets of effectors in sequential waves at different phases of parasitism. Pathogens have convergently evolved different ways to achieve this temporal regulation. At this point, these processes are better understood for animal parasites. During infection of vertebrate cells, the protozoan pathogen *Toxoplasma gondii* secretes effectors by sequential triggering of three distinct parasite organelles known as micronemes, rhoptries, and dense granules, at different phases during its intracellular growth (Carruthers and Sibley 1997). Bacterial enteropathogen *Salmonella enterica* uses a sorting platform involving chaperone proteins to sort and load the effectors to be secreted in a defined order via T3SS (Lara-Tejero et al. 2011). In bacterial plant pathogens, T3SS is also used to deliver effectors into plant cells in a regulated manner. T3SS effectors released early in the epiphytic growth of *P. syringae* *PsyB728a* are important for pathogen survival during this phase and T3SS effectors such as HopAA1 and HopZ3 may play specific roles during epiphytic growth of this pathogen on specific hosts (Lee et al. 2012b). Phytoplasmas, which require insect vectors for transmission to plants, differentially regulate effector genes for insect and plant colonization (Toruno et al. 2010; MacLean et al. 2011). Fungal pathogens *Colletotrichum higginsianum* and *Colletotrichum graminicola*, infecting *Arabidopsis thaliana* and maize, respectively, use transcriptional regulation to synthesize and secrete different sets of effectors and enzymes important for different phases of infection: The biotrophic phase is augmented by effectors and secondary metabolism enzymes whereas the necrotrophic phase brings in hydrolases and transporters (O'Connell et al. 2012). In *C. higginsianum*, the regulation is further refined to the prepenetration stage where ChCEs (*C. higginsianum* candidate effectors) are up-regulated presumably in preparation for host-cell ingress (Kleemann et al. 2012). In the oomycete *Phytophthora*, several RXLR-type effectors are induced at the cyst germination stage or at early infection stage, whereas several necrosis-inducing Nep1-like proteins (NLPs) are expressed at late infection stages, suggesting that they contribute to the necrotrophic phase (Qutob et al. 2002; Haas et al. 2009). In *P. infestans*, an effector known as SNE1 is transcriptionally regulated and expressed during the biotrophic phase, presumably to maintain biotrophy and to suppress the actions of a cell-death-inducing NLP that is expressed during the necrotrophic phase of infection (Kelley et al. 2010). Once the effectors reach the right location in host cells at the right time, they can interact with host proteins to exert their functions for successful invasion.

HOST-CELL TARGETS OF EFFECTORS

Effectors have evolved to bind and modify “target” host molecules to perturb plant processes (Figs. 1A and 2A). Essentially, any effector activity that would increase the fitness of the microbe, its ability to colonize the host plant, and spread to other hosts could potentially evolve. Some effectors, particularly bacterial T3SS effectors, are enzymes that biochemically modify host molecules, typically impeding their function or eliminating them (Cunac et al. 2009; Deslandes and Rivas 2012). The enzymatic activities of effectors are diverse and include protease, hydrolase, phosphatase, kinase, transferase, and ubiquitin ligase activities (Shao et al. 2003; Abramovitch et al. 2006b; Janjusevic et al. 2006; Fu et al. 2007; Lee et al. 2012a; Rodriguez-Herva et al. 2012; van Damme et al. 2012). Other effectors do not carry enzymatic activities and act by binding host proteins to modulate their

Figure 2. Effectors and their host-cell helpers and targets. This genetic model describes the position of pathogen effectors and their respective host-cell helpers and targets in the signaling pathways leading to susceptibility (A) and resistance (B). Effector targets and effector helpers are distinct plant susceptibility factors. Pathogen effectors recruit host helper proteins and interact with them for proper function. Activated effectors bind cognate targets, manipulate them, and form active effector–target complexes. In a susceptible interaction, the effector–target complex is not recognized and results in an altered cellular state of effector-triggered susceptibility (ETS). In a resistant interaction, this complex triggers host recognition by cognate immune receptors leading to effector-triggered immunity (ETI).

functions. Many such effectors inhibit plant enzymes such as kinases, proteases, glucanases, and peroxidases (Tian et al. 2004, 2007; Rooney et al. 2005; Damasceno et al. 2008; Xiang et al. 2008; Song et al. 2009; Hemetsberger et al. 2012). Another group of effectors have evolved to bind nucleic acids acting as modulators of gene expression. *Xanthomonas* TAL effectors directly bind elements in plant gene promoters to activate gene expression of host genes that benefit the pathogen (Duan et al. 1999; Boch et al. 2009; Domingues et al. 2010; de Souza et al. 2012). Among the host susceptibility genes that are induced by TAL effectors are the SWEET sugar transporters that are thought to release sugar to contribute to the nutrition of the invading bacteria (Chen et al. 2010).

Our knowledge of the identity of the host targets of effectors remains incomplete given the large number of uncharacterized effectors from a multitude of pathogens and symbionts. The full range of biochemical mechanisms by which effectors manipulate their targets is also poorly known. Initially, effectors were thought to be quite specific, targeting and modifying individual plant proteins. It now appears that effectors can be promiscuous; a single effector can associate with multiple plant proteins and may even affect distinct processes in the host plant (Fig. 2A). Classic examples include the *P. syringae* type-III effectors AvrPto and AvrPtoB that directly bind and interfere with several immune receptor kinases in tomato and *Arabidopsis* thereby interfering with multiple PTI signaling pathways (Abramovitch et al. 2006a; Göhre et al. 2008; Shan et al. 2008; Xiang et al. 2008; Gimenez-Ibanez et al. 2009; Zeng et al. 2012) and HopM1, which was shown by Nomura et al. (2006) to degrade a number of plant proteins in addition to AtMin7. Similarly, phytoplasma protein effector SAP11 binds to both plant class I and II TCP transcription factors but only destabilizes the latter (Sugio et al. 2011).

HOST-CELL HELPERS OF EFFECTORS

Not all effector-associated plant proteins are host targets that are modified by the effectors. Some plant proteins biochemically modify the effectors, contribute to effector maturation inside the plant cytoplasm, or serve as cofactors that form biochemically active complexes with the effector. Other host proteins enable the effectors to traffic to their final subcellular destination. Such proteins are best viewed as “effector helpers” rather than genuine “effector targets.” These effector-associated helper proteins can be distinguished from effector targets because they genetically function upstream of the effector-mediated target modulation (Fig. 2B). Loss-of-function mutations in effector helpers typically interfere with the ability of the effector to perform its activity but do not necessarily affect the biological process targeted by the effector. In contrast, mutations in effector targets usually phenocopy the phenotype induced by the effector, such as suppression of host immunity or altered plant development (Block et al. 2008; Block and Alfano 2011; Canonne and Rivas 2012; Deslandes and Rivas 2012).

There are several vivid examples of host effector helpers. The type-III effector AvrRpt2 is a cysteine protease that is delivered inside plant cells by *P. syringae* as an inactive enzyme. Once inside the host cell, AvrRpt2 requires cyclophilin, a chaperone that catalyzes and accelerates protein folding (Coaker et al. 2005, 2006). Cyclophilin acts as a cofactor, activating the self-processing of AvrRpt2 and leading to the cleavage of AvrRpt2 target protein RIN4 (Coaker et al. 2005). Similarly, another *P. syringae* type-III effector, HopZ1a, is activated inside the host cell by phytic acid to become a functional acetyltransferase that acetylates tubulin, a plant target of HopZ1a (Lee et al. 2012a). Other type-III effectors require host-mediated biochemical modifications, such as myristoylation and phosphorylation, to become functional. For example, myristoylation of AvrRpm1 and AvrPto is required for both their avirulence and virulence activities, suggesting that plant N-myristoyltransferases are involved as host-cell helpers (Nimchuk et al. 2000; Anderson et al. 2006).

Another example of a host-cell helper is the plant protein importin- α that mediates nuclear trafficking and is recruited by several pathogen effectors including TAL effectors, SAP11, and Crinklers (Szurek et al. 2001; Bai et al. 2009; Schornack et al. 2010). TAL effector AvrBs3 in association with importin- α enables its accumulation in the host-cell nucleus where it directly binds to plant DNA harboring a conserved promoter element, which is its “real” target, to activate plant gene expression (Szurek et al. 2001; Kay et al. 2007, 2009). TAL effectors also likely associate with components of the plant transcriptional machinery. One candidate is the general transcription factor subunit TFIIA γ , which is encoded by the rice recessive *Xanthomonas* resistance gene *xa5* and may serve as a cofactor that enables TAL effectors to induce host gene expression (Iyer and McCouch 2004).

Effector helpers and targets are host proteins that are usurped or manipulated by the invading organism to enable the effectors to function and establish a state of effector-triggered host susceptibility. Hence, collectively they are both host susceptibility factors. However, the distinction between helpers and targets is conceptually useful for a mechanistic dissection of effector function, particularly because mutations in these two classes of host susceptibility factors yield different phenotypes.

EFFECTOR-TARGETED PATHWAYS: FUNCTIONAL REDUNDANCY AMONG PATHOGEN EFFECTORS

Pathogens must perturb key plant processes to establish a state of susceptibility in their host cells. Interestingly, a given pathogen taxon tends to converge on particular ETPs (Fig. 3). RNA silencing (RNAi) is an efficient antiviral defense system in plants. Consequently, successful plant viruses evolved effectors that suppress RNA silencing (Burgan and Havelda 2011). Bacterial plant pathogens need to combat plant immunity elicited by conserved patterns such as flagellin and the elongation

Figure 3. Functional redundancy among effectors. This scheme illustrates how effectors target different steps in an ETP or converge on a single target within an ETP. Several unrelated effectors can also converge onto the same host protein or hub (boldface arrow) in the ETP. This reinforces the strategy of invasive pathogens to interfere and circumvent generated host signals and defense.

factor thermo unstable (EF-Tu). Therefore, many of the bacterial type-III effectors suppress plant immunity, particularly the signaling pathways triggered by flagellin and EF-Tu (Block and Alfano 2011). Similarly, fungal plant pathogens have evolved effectors that disable the immune pathway triggered by the cell wall molecule chitin (de Jonge et al. 2011). Because the signaling pathways of the immune receptors detecting flagellin, EF-Tu, and chitin overlap, it is possible that fungal effectors can target the pathways activated by bacterial flagellin and EF-Tu and vice versa. Diverse phytoplasma species elicit developmental malformations in their host plants (Sugio and Hogenhout 2012). They secrete effectors that modulate plant development and plant–insect interactions, therefore ensuring the successful transmission of these obligate and insect-vectored bacterial pathogens.

Because a given pathogen taxon must perturb specific host pathways to be effective, it is perhaps not surprising that many effectors appear to be functionally redundant. As schematically depicted in Figure 3, functionally redundant effectors may affect different steps in an ETP or may converge on a single target within the pathway. For example, three different *P. syringae* type-III effectors bind and perturb the same *Arabidopsis* target protein RIN4 (Grant et al. 2006). Two of these effectors, AvrRpm1 and AvrB, induce phosphorylation of RIN4, whereas the protease AvrRpt2 cleaves RIN4. Two additional *P. syringae* type-III effectors, AvrPto and AvrPtoB, physically interact with the kinase domains of the im-

munoreceptors FLS2, EFR, and CERK1, thereby interfering with multiple PTI signaling pathways (Nicaise et al. 2009). In contrast, the effectors HopA11 and HopF2 inhibit downstream steps of the same PTI signaling pathways targeted by AvrPto and AvrPtoB. HopA11 and HopF2 target MPKs and MKKs, respectively, which are kinases that mediate PTI signaling following perception of microbial patterns (Wang et al. 2010; Wu et al. 2011). This example also illustrates how the same effector could target two or more different steps within an ETP (Fig. 3).

The effectors of filamentous pathogens also display functional redundancy. Both of the effectors Avr4 and Ecp6 of the leaf mold fungus *C. fulvum* interfere with chitin perception by the host plant tomato (de Jonge et al. 2011). Likewise, the RXLR-type effectors of oomycetes are thought to exhibit a high degree of functional redundancy. Genome and transcriptome analyses of different strains of the potato late blight pathogen *Phytophthora infestans* revealed a high degree of genetic and expression polymorphism (Haas et al. 2009; Raffaele et al. 2010; Cooke et al. 2012). Many effector genes are strain specific: Only 45 of the ~550 predicted RXLR effector genes have intact coding sequences and are induced in planta in all examined strains (Cooke et al. 2012). In the soybean pathogen *Phytophthora sojae*, 10%–15% of RXLR effectors are highly expressed during infection (Jiang and Tyler 2012). This suggests that the majority of the RXLR effectors of *P. infestans* and *P. sojae* are dispensable and possibly encode redundant functions.

Effector functional redundancy may also be a consequence of arms race coevolution between pathogens and hosts. The pathogen population could benefit from carrying a set of functionally redundant effectors to counteract plant immunity. Genetic and epigenetic modifications of gene function, including nonsynonymous mutations, deletion, and gene silencing, are usually deleterious, being removed from the pathogen population by purifying selection. On the other hand, purifying selection is unlikely to effectively operate on redundant effector genes because they functionally complement one another. This results in the maintenance of genes with relatively low fitness under the present environmental conditions and facilitates the emergence of strain-specific repertoires of effector genes in the population. When the environment changes, for instance, through the emergence of disease resistance in the host plant, preexisting isolates harboring a set of effectors advantageous for the new conditions could be positively selected. This evolutionary “bet-hedging” strategy, in which the pathogen population benefits from extensive heterogeneity in the effector complement, could be one of the tactics that enable pathogens to avoid extinction in the arms race (Toruno et al. 2010; Simons 2011). For example, a functionally redundant heterogeneous effector repertoire would enable the pathogen population to include strains that can disable key ETPs while evading perception by plant immunoreceptors (Boller and Felix 2009). The extent to which “bet hedging” explains functional redundancy of plant pathogen effectors has not been explored

in depth and deserves to be more rigorously studied in the future.

EFFECTORS FROM UNRELATED PATHOGENS CONVERGE ON THE SAME HOST TARGETS

We discussed above how different effectors from the same pathogen have evolved to disable the same host target within an ETP (Fig. 3). In addition, effectors from phylogenetically unrelated pathogens can also converge on the same host targets. One example is the tomato defense cysteine protease RCR3, which is inhibited by effectors from three unrelated plant pathogens: *C. fulvum* AVR2, *P. infestans* EPIC1 and EPIC2B, and VAP1 from the root parasitic nematode *Globodera rostochiensis* (Song et al. 2009; Lozano-Torres et al. 2012). These findings highlight RCR3 as a “hub” component of apoplastic defenses of tomato that needs to be disabled for successful parasitism by fungal, oomycete, and nematode pathogens. Remarkably, plants have evolved to detect pathogen-induced perturbations of hub virulence targets such as RCR3. The tomato cell surface receptor-like protein Cf-2 confers resistance to *C. fulvum* and *G. rostochiensis* in an RCR3-dependent manner (see more on this topic in the following section) (Rooney et al. 2005; Lozano-Torres et al. 2012). More recently, Mukhtar et al. (2011) identified a set of 18 candidate hub target proteins following a large-scale yeast two-hybrid (Y2H) screen using effectors of both the oomycete *Hyaloperonospora arabidopsidis* and the bacterium *P. syringae*.

Although a number of effectors from unrelated pathogens have evolved to disable identical targets, in many cases any step in an ETP would be an effective target. Is it truly more beneficial for a pathogen to converge on a few “hub” host targets given that nodes would then be under strong selection to evade effectors? It seems that in many cases, pathogens have evolved effectors that disable multiple steps within a targeted pathway. For instance, plant viruses have evolved a multitude of mechanisms to target the RNA silencing machinery but have not evolved to focus on a single hub target (Burgyn and Havelda 2011).

Perhaps a more pertinent issue is to determine the degree to which unrelated pathogen taxa have converged on the same ETPs. Some common themes are starting to emerge. Effectors from diverse pathogenic species of bacteria, oomycetes, fungi, and insects have been reported to suppress an overlapping set of PTI signaling pathways (Gimenez-Ibanez et al. 2009; Bos et al. 2010; de Jonge et al. 2010; Wang et al. 2011). Both bacterial and oomycete plant pathogens target vesicle trafficking pathways to interfere with focal immunity, although the exact mechanisms are yet to emerge (Bozkurt et al. 2012; Lindeberg et al. 2012). As discussed above, filamentous pathogens and nematodes have adapted to the proteolytic environment of the plant apoplast by evolving diverse protease inhibitor effectors (Rooney et al. 2005; Song et al. 2009; Lozano-Torres et al. 2012).

ETPs are a direct reflection of the pathogenic strategy used by a given pathogen and the obstacles that the pathogen faces during host infection. In the future, as we discover ETPs for a wider range of plant-interacting taxa, we should be able to classify and define plant pathogens and pests by the nature of the host processes that they must impair.

RECOGNITION OF PATHOGEN EFFECTORS BY PLANT IMMUNORECEPTORS

Plants have evolved a potent immune system to recognize invading organisms. This ubiquitous immune response uses conserved pathways to respond to a variety of parasites, including viruses, bacteria, fungi, oomycetes, nematodes, insects, and parasitic plants. Plant immunoreceptors can turn the table on the parasite by recognizing effector molecules, therefore turning the effector into a liability for the pathogen (Figs. 1B and 2B).

Plant perception of invading organisms can occur via cell surface receptors, receptor-like kinases or receptor-like proteins, or NB-LRR proteins, a hugely diverse class of plant intracellular immunoreceptors. These immune receptors can perceive all types of pathogen molecules, ranging from conserved microbial patterns (also known as PAMPs) to highly polymorphic and rapidly evolving effectors. Recognition of pathogen molecules can be direct or indirect. The cell surface PRRs FLS2 and EFR directly bind their ligands to the PAMPs flagellin and EF-Tu, respectively (Chinchilla et al. 2006; Zipfel et al. 2006). Another example of direct recognition is the direct binding between the flax NB-LRR receptors L5, L6, and L7 and the flax rust fungus AvrL567 effector (Dodds et al. 2006). In contrast, in plant–bacteria interactions, intracellular immunoreceptors often recognize effectors indirectly through the perception of modifications in the host protein targeted by the effector. For instance, *Arabidopsis* NB-LRR protein RPM1 detects modifications in the host protein RIN4, which is targeted by the *P. syringae* effector proteins AvrB and AvrRpm1 (Mackey et al. 2002). With no universal recognition mechanism evident, several key models have been proposed to encapsulate the complexity of the recognition process.

Historically, indirect recognition was viewed as a guard process, with immunoreceptors acting as sentinels that “guard” particular plant proteins and detect pathogen-induced perturbations (Jones and Dangl 2006). This model had to be expanded with the discovery that most effectors bind multiple host targets, some of which having no apparent role in ETS. One theory is that the plant deploys decoy proteins to sequester the effectors and activate innate immunity. In the decoy model, decoy proteins act exclusively in effector recognition, mimicking the operative target (van der Hoorn and Kamoun 2008). For instance, indirect recognition of Avr2 by tomato Cf-2 is mediated by the selective inhibition of two papain-like cysteine proteases, RCR3 and PIP1, by Avr2 (Rooney et al. 2005; Shabab et al. 2008). Shabab et al. (2008)

proposed that PIP1 is the operative target of Avr2 and that the decoy RCR3 acts to exclusively mediate recognition by Cf-2. Accordingly, RCR3 accumulates in the tomato apoplast at much lower levels than PIP1 during infection, and inhibition of RCR3 does not contribute to enhanced virulence (Dixon et al. 2000).

In addition to models involving NB-LRR immunoreceptors, the recognition of bacterial TAL effectors can present a distinct mechanism. Some resistance genes that recognize TAL effectors are transcriptionally activated through the interaction of corresponding effectors with elements within their promoters (see above). For example, the pepper resistance gene *Bs3* is targeted by the *Xanthomonas* AvrBs3 effector that directly activates its transcription and thereby confers resistance (Romer et al. 2007).

EFFECTORS AS MOLECULAR PROBES TO STUDY PLANT BIOLOGY

Research on plant–microbe interactions has yielded many key discoveries that impacted all of plant biology. For example, the plant hormone gibberellin was first identified as the molecule that triggers the “foolish rice” symptoms caused by the plant pathogenic fungus *Gibberella fujikuroi* (Tamura 1990). Strigolactones, which are now known to inhibit plant shoot branching, were first discovered as stimulants of seed germination of parasitic plants and growth activators of arbuscular mycorrhizal fungi (Xie and Yoneyama 2010). Other areas of plant biology that have been significantly influenced by the study of plant–biotic interactions include developmental biology, receptor biology, and epigenetics (Speth et al. 2007).

The study of the effectors of plant-associated organisms promises to continue this tradition of influential and wide-reaching discoveries. Effectors can be used as molecular probes to unravel unknown aspects of plant immunity and biology (Bozkurt et al. 2012; Feng and Zhou 2012). By definition, studying the function of an effector is a leap into the unknown. One has little clue regarding where the research will lead. But the targeted host process must be important enough to enable the effector activity to emerge and become selected. In essence, effectors have evolved as a result of an awesome, comprehensive genetic screen that has taken place in nature over millions of years.

Effectors have been used as molecular probes in multiple ways. The type-III effectors of pathogenic bacteria turned out to be an exquisite toolkit to dissect plant immune pathways (Feng and Zhou 2012; Wei et al. 2012). Effectors are also useful probes for plant cell biology. Plant vesicular trafficking and focal immunity have been difficult processes to dissect using standard genetic approaches, mainly because mutants tend to be pleiotropic. Effectors promise to be a useful toolkit to dissect such dynamic cellular processes, potentially helping to unravel the diversity of secretory vesicles and their cargo (Bozkurt et al. 2011).

SYNTHETIC EFFECTORS: NEW TOOLS FOR PLANT BIOTECHNOLOGY

Plant pathogen effectors have proven to be a rich source of genetic innovation, often evolving independent mechanisms that mimic plant processes. However, effectors have evolved to benefit the invading organism. Can we turn this around and engineer beneficial synthetic effectors that can be used in biotechnology? This has already happened with TAL effectors, which can be custom designed and engineered to bind any target DNA sequence (Bogdanove and Voytas 2011). Indeed, TAL effectors have emerged as key tools for synthetic biology. TAL effectors can be fused to DNA nucleases to target a precise site in a genome to produce genetic variations. Such TAL nucleases have already been used to produce targeted variations in the genomes of mammals, flies, worms, and plants (Bogdanove and Voytas 2011). In a recent landmark study, Li et al. (2012) used TAL nucleases to generate 3–57 base deletions in a specific susceptibility gene of rice to engineer bacterial blight resistance. Other type-III effectors can also be used as reagents to manipulate cellular pathways. Wei et al. (2012) engineered two type-III effector proteins from animal parasitic bacteria to rewire kinase pathways in yeast and mammalian cells. In the future, as we learn more about structure–function relationships and mechanisms of action of different classes of effectors, we can further exploit effectors as synthetic biology reagents.

CONCLUDING REMARKS

The effectors of plant-associated organisms are a source of biological innovation; they include some of the most remarkable proteins known to function in plant cells and tissue. Nonetheless, the field of effector biology remains in its infancy. Our knowledge of effectors is mostly limited to a few taxa of plant pathogens and to immunosuppression processes. We know little about the identity and function of effectors from deep lineages of plant-associated organisms, including obligate plant pathogens, such as powdery mildews and rust fungi, and mutualists, such as mycorrhizal fungi. In addition, the occurrence of effectors in insect herbivores and parasitic plants is only starting to be appreciated (Gheysen and Mitchum 2011; Hogenhout and Bos 2011). Clearly, this plethora of effectors from a wide range of plant-associated organisms are likely to have unexpected functions and should prove to be a rich source of scientific discovery. Effector biology is, therefore, poised to continue to broadly impact the biological sciences.

ACKNOWLEDGMENTS

The ideas discussed in this article were shaped by discussions with many students and colleagues. We thank the participants of the 2012 NOVA Ph.D. course in plant pathology, the EMBO course “Plant-Microbe Interactions,” particularly, and Isabel Weinheime and Benjamin Petre for their contributions. We are also grateful to the

tweeps of the @KamounLab Twitter account for useful discussions. Finally, we acknowledge funding from the Biotechnology and Biological Sciences Research Council (BBSRC), the European Research Council (ERC), the Leverhulme Trust Early Career Fellowship to D.G.O.S., a Japan Society for the Promotion of Science Postdoctoral Fellowship for Research Abroad to K.Y., and the Gatsby Charitable Foundation.

REFERENCES

- Abramovitch RB, Anderson JC, Martin GB. 2006a. Bacterial elicitation and evasion of plant innate immunity. *Nat Rev Mol Cell Biol* **7**: 601–611.
- Abramovitch RB, Janjusevic R, Stebbins CE, Martin GB. 2006b. Type III effector AvrPtoB requires intrinsic E3 ubiquitin ligase activity to suppress plant cell death and immunity. *Proc Natl Acad Sci* **103**: 2851–2856.
- Agrios GN. 2005. *Plant Pathology*, 5th ed. Elsevier, San Diego.
- Anderson JC, Pascuzzi PE, Xiao F, Sessa G, Martin GB. 2006. Host-mediated phosphorylation of type III effector AvrPto promotes *Pseudomonas* virulence and avirulence in tomato. *Plant Cell* **18**: 502–514.
- Bai X, Correa VR, Toruno TY, Ammar el D, Kamoun S, Hogenhout SA. 2009. AY-WB phytoplasma secretes a protein that targets plant cell nuclei. *Mol Plant Microbe Interact* **22**: 18–30.
- Bartetzko V, Sonnewald S, Vogel F, Hartner K, Stadler R, Hammes UZ, Bornke F. 2009. The *Xanthomonas campestris* pv *vesicatoria* type III effector protein XopJ inhibits protein secretion: Evidence for interference with cell wall-associated defense responses. *Mol Plant Microbe Interact* **22**: 655–664.
- Bender CL, Alarcon-Chaidez F, Gross DC. 1999. *Pseudomonas syringae* phytotoxins: Mode of action, regulation, and biosynthesis by peptide and polyketide synthetases. *Microbiol Mol Biol Rev* **63**: 266–292.
- Block A, Alfano JR. 2011. Plant targets for *Pseudomonas syringae* type III effectors: Virulence targets or guarded decoys? *Curr Opin Microbiol* **14**: 39–46.
- Block A, Li G, Fu ZQ, Alfano JR. 2008. Phytopathogen type III effector weaponry and their plant targets. *Curr Opin Plant Biol* **11**: 396–403.
- Block A, Guo M, Li G, Elowsky C, Clemente TE, Alfano JR. 2010. The *Pseudomonas syringae* type III effector HopG1 targets mitochondria, alters plant development and suppresses plant innate immunity. *Cell Microbiol* **12**: 318–330.
- Boch J, Scholze H, Schornack S, Landgraf A, Hahn S, Kay S, Lahaye T, Nickstadt A, Bonas U. 2009. Breaking the code of DNA binding specificity of TAL-type III effectors. *Science* **326**: 1509–1512.
- Bogdanove AJ, Voytas DF. 2011. TAL effectors: Customizable proteins for DNA targeting. *Science* **333**: 1843–1846.
- Boller T, Felix G. 2009. A renaissance of elicitors: Perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu Rev Plant Biol* **60**: 379–406.
- Bolton MD, van Esse HP, Vossen JH, de Jonge R, Stergiopoulos I, Stulemeijer IJ, van den Berg GC, Borrás-Hidalgo O, Dekker HL, de Koster CG, et al. 2008. The novel *Cladosporium fulvum* lysin motif effector Ecp6 is a virulence factor with orthologues in other fungal species. *Mol Microbiol* **69**: 119–136.
- Bos JI, Prince D, Pitino M, Maffei ME, Win J, Hogenhout SA. 2010. A functional genomics approach identifies candidate effectors from the aphid species *Myzus persicae* (green peach aphid). *PLoS Genet* **6**: e1001216. doi: 10.1371/journal.pgen.1001216.
- Bozkurt TO, Schornack S, Win J, Shindo T, Ilyas M, Oliva R, Cano LM, Jones AM, Huitema E, van der Hoorn RA, et al. 2011. *Phytophthora infestans* effector AVRblb2 prevents secretion of a plant immune protease at the haustorial interface. *Proc Natl Acad Sci* **108**: 20832–20837.
- Bozkurt TO, Schornack S, Banfield MJ, Kamoun S. 2012. Oomycetes, effectors, and all that jazz. *Curr Opin Plant Biol* **15**: 483–492.
- Burgan J, Havelda Z. 2011. Viral suppressors of RNA silencing. *Trends Plant Sci* **16**: 265–272.
- Buttner D, He SY. 2009. Type III protein secretion in plant pathogenic bacteria. *Plant Physiol* **150**: 1656–1664.
- Caillaud M-C, Wirthmueller L, Fabro G, Piquerez SJM, Asai S, Ishaque N, Jones JDG. 2012. Mechanisms of nuclear suppression of host immunity by effectors from the *Arabidopsis* downy mildew pathogen *Hyaloperonospora arabidopsidis* (Hpa). *Cold Spring Harb Symp Quant Biol* **77**: doi: 10.1101/sqb.2012.77.a015115.
- Canonne J, Rivas S. 2012. Bacterial effectors target the plant cell nucleus to subvert host transcription. *Plant Signal Behav* **7**: 217–221.
- Carruthers VB, Sibley LD. 1997. Sequential protein secretion from three distinct organelles of *Toxoplasma gondii* accompanies invasion of human fibroblasts. *Eur J Cell Biol* **73**: 114–123.
- Chen LQ, Hou BH, Lalonde S, Takanaga H, Hartung ML, Qu XQ, Guo WJ, Kim JG, Underwood W, Chaudhuri B, et al. 2010. Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* **468**: 527–532.
- Chinchilla D, Bauer Z, Regenass M, Boller T, Felix G. 2006. The *Arabidopsis* receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception. *Plant Cell* **18**: 465–476.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ. 2006. Host-microbe interactions: Shaping the evolution of the plant immune response. *Cell* **124**: 803–814.
- Christie PJ. 2004. Type IV secretion: The *Agrobacterium* VirB/D4 and related conjugation systems. *Biochim Biophys Acta* **1694**: 219–234.
- Coaker G, Falick A, Staskawicz B. 2005. Activation of a phytopathogenic bacterial effector protein by a eukaryotic cyclophilin. *Science* **308**: 548–550.
- Coaker G, Zhu G, Ding Z, Van Doren SR, Staskawicz B. 2006. Eukaryotic cyclophilin as a molecular switch for effector activation. *Mol Microbiol* **61**: 1485–1496.
- Cooke DEL, Cano LM, Raffaele S, Bain RA, Cooke LR, Etherington GJ, Deahl KL, Farrer RA, Gilroy EM, Goss EM, et al. 2012. Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathog* (in press).
- Cunnac S, Lindeberg M, Collmer A. 2009. *Pseudomonas syringae* type III secretion system effectors: Repertoires in search of functions. *Curr Opin Microbiol* **12**: 53–60.
- Damasceno CM, Bishop JG, Ripoll DR, Win J, Kamoun S, Rose JK. 2008. Structure of the glucanase inhibitor protein (GIP) family from *Phytophthora* species suggests coevolution with plant endo- β -1,3-glucanases. *Mol Plant Microbe Interact* **21**: 820–830.
- de Jonge R, van Esse HP, Kombrink A, Shinya T, Desaki Y, Bours R, van der Krol S, Shibuya N, Joosten MH, Thomma BP. 2010. Conserved fungal LysM effector Ecp6 prevents chitin-triggered immunity in plants. *Science* **329**: 953–955.
- de Jonge R, Bolton MD, Thomma BP. 2011. How filamentous pathogens co-opt plants: The ins and outs of fungal effectors. *Curr Opin Plant Biol* **14**: 400–406.
- Deslandes L, Rivas S. 2012. Catch me if you can: Bacterial effectors and plant targets. *Trends Plant Sci* **17**: 644–655.
- de Souza TA, Soprano AS, de Lira NP, Quaresma AJ, Pauletti BA, Paes Leme AF, Benedetti CE. 2012. The TAL effector PthA4 interacts with nuclear factors involved in RNA-dependent processes including a HMG protein that selectively binds poly(U) RNA. *PLoS One* **7**: e32305. doi: p10.1371/journal.pone.0032305.
- Dixon MS, Golstein C, Thomas CM, van Der Biezen EA, Jones JD. 2000. Genetic complexity of pathogen perception by plants: The example of *Rcr3*, a tomato gene required specifically by *Cf-2*. *Proc Natl Acad Sci* **97**: 8807–8814.

- Dodds PN, Rathjen JP. 2010. Plant immunity: Towards an integrated view of plant-pathogen interactions. *Nat Rev Genet* **11**: 539–548.
- Dodds PN, Lawrence GJ, Catanzariti AM, Teh T, Wang CI, Ayliffe MA, Kobe B, Ellis JG. 2006. Direct protein interaction underlies gene-for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. *Proc Natl Acad Sci* **103**: 8888–8893.
- Dodds PN, Rafiqi M, Gan PH, Hardham AR, Jones DA, Ellis JG. 2009. Effectors of biotrophic fungi and oomycetes: Pathogenicity factors and triggers of host resistance. *New Phytol* **183**: 993–1000.
- Domingues MN, De Souza TA, Cernadas RA, de Oliveira ML, Docena C, Farah CS, Benedetti CE. 2010. The *Xanthomonas citri* effector protein PthA interacts with citrus proteins involved in nuclear transport, protein folding and ubiquitination associated with DNA repair. *Mol Plant Pathol* **11**: 663–675.
- Duan YP, Castaneda A, Zhao G, Erdos G, Gabriel DW. 1999. Expression of a single, host-specific, bacterial pathogenicity gene in plant cells elicits division, enlargement, and cell death. *Mol Plant Microbe Interact* **12**: 556–560.
- Feng F, Zhou JM. 2012. Plant-bacterial pathogen interactions mediated by type III effectors. *Curr Opin Plant Biol* **15**: 469–476.
- Fu ZQ, Guo M, Jeong BR, Tian F, Elthon TE, Cerny RL, Staiger D, Alfano JR. 2007. A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity. *Nature* **447**: 284–288.
- Gheysen G, Mitchum MG. 2011. How nematodes manipulate plant development pathways for infection. *Curr Opin Plant Biol* **14**: 415–421.
- Gimenez-Ibanez S, Hann DR, Ntoukakis V, Petutschnig E, Lipka V, Rathjen JP. 2009. AvrPtoB targets the LysM receptor kinase CERK1 to promote bacterial virulence on plants. *Curr Biol* **19**: 423–429.
- Göhre V, Spallek T, Haweker H, Mersmann S, Mentzel T, Boller T, de Torres M, Mansfield JW, Robatzek S. 2008. Plant pattern-recognition receptor FLS2 is directed for degradation by the bacterial ubiquitin ligase AvrPtoB. *Curr Biol* **18**: 1824–1832.
- Grant SR, Fisher EJ, Chang JH, Mole BM, Dangl JL. 2006. Subterfuge and manipulation: Type III effector proteins of phytopathogenic bacteria. *Annu Rev Microbiol* **60**: 425–449.
- Haas BJ, Kamoun S, Zody MC, Jiang RH, Handsaker RE, Cano LM, Grabherr M, Kodira CD, Raffaele S, Torto-Alalibo T, et al. 2009. Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* **461**: 393–398.
- Hemetsberger C, Herrberger C, Zechmann B, Hillmer M, Doehlemann G. 2012. The *Ustilago maydis* effector Pep1 suppresses plant immunity by inhibition of host peroxidase activity. *PLoS Pathog* **8**: e1002684. doi: 10.1371/journal.ppat.1002684.
- Hines PJ, Zahn LM. 2009. What's bugging plants? *Science* **324**: 741.
- Hogenhout SA, Bos JJ. 2011. Effector proteins that modulate plant-insect interactions. *Curr Opin Plant Biol* **14**: 422–428.
- Hogenhout SA, Loria R. 2008. Virulence mechanisms of Gram-positive plant pathogenic bacteria. *Curr Opin Plant Biol* **11**: 449–456.
- Hogenhout SA, Van der Hoorn RA, Terauchi R, Kamoun S. 2009. Emerging concepts in effector biology of plant-associated organisms. *Mol Plant Microbe Interact* **22**: 115–122.
- Iyer AS, McCouch SR. 2004. The rice bacterial blight resistance gene *xa5* encodes a novel form of disease resistance. *Mol Plant Microbe Interact* **17**: 1348–1354.
- Janjusevic R, Abramovitch RB, Martin GB, Stebbins CE. 2006. A bacterial inhibitor of host programmed cell death defenses is an E3 ubiquitin ligase. *Science* **311**: 222–226.
- Jelenska J, Yao N, Vinatzer BA, Wright CM, Brodsky JL, Greenberg JT. 2007. A J domain virulence effector of *Pseudomonas syringae* remodels host chloroplasts and suppresses defenses. *Curr Biol* **17**: 499–508.
- Jiang RH, Tyler BM. 2012. Mechanisms and evolution of virulence in oomycetes. *Annu Rev Phytopathol* **50**: 295–318.
- Jones JD, Dangl JL. 2006. The plant immune system. *Nature* **444**: 323–329.
- Kaffarnik FA, Jones AM, Rathjen JP, Peck SC. 2009. Effector proteins of the bacterial pathogen *Pseudomonas syringae* alter the extracellular proteome of the host plant, *Arabidopsis thaliana*. *Mol Cell Proteomics* **8**: 145–156.
- Kakizawa S, Oshima K, Nishigawa H, Jung HY, Wei W, Suzuki S, Tanaka M, Miyata S, Ugaki M, Namba S. 2004. Secretion of immunodominant membrane protein from onion yellows phytoplasma through the Sec protein-translocation system in *Escherichia coli*. *Microbiology* **150**: 135–142.
- Kay S, Hahn S, Marois E, Hause G, Bonas U. 2007. A bacterial effector acts as a plant transcription factor and induces a cell size regulator. *Science* **318**: 648–651.
- Kay S, Hahn S, Marois E, Wieduwild R, Bonas U. 2009. Detailed analysis of the DNA recognition motifs of the *Xanthomonas* type III effectors AvrBs3 and AvrBs3Deltarep16. *Plant J* **59**: 859–871.
- Kelley BS, Lee SJ, Damasceno CM, Chakravarthy S, Kim BD, Martin GB, Rose JK. 2010. A secreted effector protein (SNE1) from *Phytophthora infestans* is a broadly acting suppressor of programmed cell death. *Plant J* **62**: 357–366.
- Kleemann J, Rincon-Rivera LJ, Takahara H, Neumann U, van Themaat EV, van der Does HC, Hacquard S, Stuber K, Will I, Schmalenbach W, et al. 2012. Sequential delivery of host-induced virulence effectors by appressoria and intracellular hyphae of the phytopathogen *Colletotrichum higginsianum*. *PLoS Pathog* **8**: e1002643. doi: 10.1371/journal.ppat.1002643.
- Kloppholz S, Kuhn H, Requena N. 2011. A secreted fungal effector of *Glomus intraradices* promotes symbiotic biotrophy. *Curr Biol* **21**: 1204–1209.
- Lakatos L, Szittyá G, Silhavy D, Burgyn J. 2004. Molecular mechanism of RNA silencing suppression mediated by p19 protein of tombusviruses. *EMBO J* **23**: 876–884.
- Lara-Tejero M, Kato J, Wagner S, Liu X, Galan JE. 2011. A sorting platform determines the order of protein secretion in bacterial type III systems. *Science* **331**: 1188–1191.
- Lee AH, Hurley B, Felsensteiner C, Yea C, Ckurshumova W, Bartetzko V, Wang PW, Quach V, Lewis JD, Liu YC, et al. 2012a. A bacterial acetyltransferase destroys plant microtubule networks and blocks secretion. *PLoS Pathog* **8**: e1002523. doi: 10.1371/journal.ppat.1002523.
- Lee J, Teitzel GM, Munkvold K, del Pozo O, Martin GB, Michelmore RW, Greenberg JT. 2012b. Type III secretion and effectors shape the survival and growth pattern of *Pseudomonas syringae* on leaf surfaces. *Plant Physiol* **158**: 1803–1818.
- Li T, Liu B, Spalding MH, Weeks DP, Yang B. 2012. High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nat Biotechnol* **30**: 390–392.
- Lindeberg M, Cunnac S, Collmer A. 2012. *Pseudomonas syringae* type III effector repertoires: Last words in endless arguments. *Trends Microbiol* **20**: 199–208.
- Lorang JM, Sweat TA, Wolpert TJ. 2007. Plant disease susceptibility conferred by a “resistance” gene. *Proc Natl Acad Sci* **104**: 14861–14866.
- Lozano-Torres JL, Wilbers RH, Gawronski P, Boshoven JC, Finkers-Tomczak A, Cordewener JH, America AH, Overmars HA, Van't Klooster JW, Baranowski L, et al. 2012. Dual disease resistance mediated by the immune receptor Cf-2 in tomato requires a common virulence target of a fungus and a nematode. *Proc Natl Acad Sci* **109**: 10119–10124.
- Mackey D, Holt BF, Wiig A, Dangl JL. 2002. RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* **108**: 743–754.
- MacLean AM, Sugio A, Makarova OV, Findlay KC, Grieve VM, Toth R, Nicolaisen M, Hogenhout SA. 2011. Phytoplasma effector SAP54 induces indeterminate leaf-like flower development in *Arabidopsis* plants. *Plant Physiol* **157**: 831–841.

- Mengiste T. 2012. Plant immunity to necrotrophs. *Annu Rev Phytopathol* **50**: 267–294.
- Mukhtar MS, Carvunis AR, Dreze M, Epple P, Steinbrenner J, Moore J, Tasan M, Galli M, Hao T, Nishimura MT, et al. 2011. Independently evolved virulence effectors converge onto hubs in a plant immune system network. *Science* **333**: 596–601.
- Nicaise V, Roux M, Zipfel C. 2009. Recent advances in PAMP-triggered immunity against bacteria: Pattern recognition receptors watch over and raise the alarm. *Plant Physiol* **150**: 1638–1647.
- Nimchuk Z, Marois E, Kjemtrup S, Leister RT, Katagiri F, Dangel JL. 2000. Eukaryotic fatty acylation drives plasma membrane targeting and enhances function of several type III effector proteins from *Pseudomonas syringae*. *Cell* **101**: 353–363.
- Nomura K, Debroy S, Lee YH, Pumplin N, Jones J, He SY. 2006. A bacterial virulence protein suppresses host innate immunity to cause plant disease. *Science* **313**: 220–223.
- Nomura K, Mecey C, Lee YN, Imboden LA, Chang JH, He SY. 2011. Effector-triggered immunity blocks pathogen degradation of an immunity-associated vesicle traffic regulator in *Arabidopsis*. *Proc Natl Acad Sci* **108**: 10774–10779.
- O'Connell RJ, Thon MR, Hacquard S, Amyotte SG, Kleemann J, Torres MF, Damm U, Buiaite EA, Epstein L, Alkan N, et al. 2012. Lifestyle transitions in plant pathogenic *Colletotrichum* fungi deciphered by genome and transcriptome analyses. *Nat Genet* **44**: 1060–1065.
- Plett JM, Kemppainen M, Kale SD, Kohler A, Legue V, Brun A, Tyler BM, Pardo AG, Martin F. 2011. A secreted effector protein of *Laccaria bicolor* is required for symbiosis development. *Curr Biol* **21**: 1197–1203.
- Qutob D, Kamoun S, Gijzen M. 2002. Expression of a *Phytophthora sojae* necrosis-inducing protein occurs during transition from biotrophy to necrotrophy. *Plant J* **32**: 361–373.
- Raffaële S, Farrer RA, Cano LM, Studholme DJ, MacLean D, Thines M, Jiang RHY, Zody MC, Kunjeti SG, Donofrio NM, et al. 2010. Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science* **330**: 1540–1543.
- Redford AJ, Bowers RM, Knight R, Linhart Y, Fierer N. 2010. The ecology of the phyllosphere: Geographic and phylogenetic variability in the distribution of bacteria on tree leaves. *Environ Microbiol* **12**: 2885–2893.
- Rodriguez RJ, White JF Jr, Arnold AE, Redman RS. 2009. Fungal endophytes: Diversity and functional roles. *New Phytol* **182**: 314–330.
- Rodriguez-Herva JJ, Gonzalez-Melendi P, Cuartas-Lanza R, Antunez-Lamas M, Rio-Alvarez I, Li Z, Lopez-Torrejón G, Diaz I, Del Pozo JC, Chakravarthy S, et al. 2012. A bacterial cysteine protease effector protein interferes with photosynthesis to suppress plant innate immune responses. *Cell Microbiol* **14**: 669–681.
- Romer P, Hahn S, Jordan T, Strauss T, Bonas U, Lahaye T. 2007. Plant pathogen recognition mediated by promoter activation of the pepper Bs3 resistance gene. *Science* **318**: 645–648.
- Rooney HC, Van't Klooster JW, van der Hoorn RA, Joosten MH, Jones JD, de Wit PJ. 2005. *Cladosporium* Avr2 inhibits tomato Rcr3 protease required for Cf-2-dependent disease resistance. *Science* **308**: 1783–1786.
- Rosebrock TR, Zeng L, Brady JJ, Abramovitch RB, Xiao F, Martin GB. 2007. A bacterial E3 ubiquitin ligase targets a host protein kinase to disrupt plant immunity. *Nature* **448**: 370–374.
- Saunders DG, Breen S, Win J, Schornack S, Hein I, Bozkurt TO, Champouret N, Vleeshouwers VG, Birch PR, Gilroy EM, et al. 2012. Host Protein BSL1 associates with *Phytophthora infestans* RXLR effector AVR2 and the *Solanum demissum* immune receptor R2 to mediate disease resistance. *Plant Cell* **24**: 3420–3434.
- Schornack S, van Damme M, Bozkurt TO, Cano LM, Smoker M, Thines M, Gaulin E, Kamoun S, Huitema E. 2010. Ancient class of translocated oomycete effectors targets the host nucleus. *Proc Natl Acad Sci* **107**: 17421–17426.
- Shabab M, Shindo T, Gu C, Kaschani F, Pansuriya T, Chintla R, Harzen A, Colby T, Kamoun S, van der Hoorn RA. 2008. Fungal effector protein AVR2 targets diversifying defense-related cysteine proteases of tomato. *Plant Cell* **20**: 1169–1183.
- Shan L, He P, Li J, Heese A, Peck SC, Nurnberger T, Martin GB, Sheen J. 2008. Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host Microbe* **4**: 17–27.
- Shao F, Golstein C, Ade J, Stoutemyer M, Dixon JE, Innes RW. 2003. Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. *Science* **301**: 1230–1233.
- Simons AM. 2011. Modes of response to environmental change and the elusive empirical evidence for bet hedging. *Proc Biol Sci* **278**: 1601–1609.
- Song J, Win J, Tian M, Schornack S, Kaschani F, Ilyas M, van der Hoorn RA, Kamoun S. 2009. Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens target the tomato defense protease Rcr3. *Proc Natl Acad Sci* **106**: 1654–1659.
- Speth EB, Lee YN, He SY. 2007. Pathogen virulence factors as molecular probes of basic plant cellular functions. *Curr Opin Plant Biol* **10**: 580–586.
- Sugio A, Hogenhout SA. 2012. The genome biology of phytoplasma: Modulators of plants and insects. *Curr Opin Microbiol* **15**: 247–254.
- Sugio A, Kingdom HN, MacLean AM, Grieve VM, Hogenhout SA. 2011. Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating plant development and defense hormone biosynthesis. *Proc Natl Acad Sci* **108**: E1254–E1263.
- Szurek B, Marois E, Bonas U, Van den Ackerveken G. 2001. Eukaryotic features of the *Xanthomonas* type III effector AvrBs3: Protein domains involved in transcriptional activation and the interaction with nuclear import receptors from pepper. *Plant J* **26**: 523–534.
- Tamura S. 1990. Historical aspects of gibberellins. In *Gibberellins* (eds Takahashi N, Phinney BO, Macmillan J), pp. 1–8. Springer-Verlag, New York.
- Tanaka H, Kitakura S, De Rycke R, De Groot R, Friml J. 2009. Fluorescence imaging-based screen identifies ARF GEF component of early endosomal trafficking. *Curr Biol* **19**: 391–397.
- Thomma BP, Nurnberger T, Joosten MH. 2011. Of PAMPs and effectors: The blurred PTI-ETI dichotomy. *Plant Cell* **23**: 4–15.
- Tian M, Huitema E, da Cunha L, Torto-Alalibo T, Kamoun S. 2004. A Kazal-like extracellular serine protease inhibitor from *Phytophthora infestans* targets the tomato pathogenesis-related protease P69B. *J Biol Chem* **279**: 26370–26377.
- Tian M, Win J, Song J, van der Hoorn R, van der Knaap E, Kamoun S. 2007. A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-like apoplastic protease. *Plant Physiol* **143**: 364–377.
- Toruno TY, Music MS, Simi S, Nicolaisen M, Hogenhout SA. 2010. Phytoplasma PMU1 exists as linear chromosomal and circular extrachromosomal elements and has enhanced expression in insect vectors compared with plant hosts. *Mol Microbiol* **77**: 1406–1415.
- Underwood W, Somerville SC. 2008. Focal accumulation of defenses at sites of fungal pathogen attack. *J Exp Bot* **59**: 3501–3508.
- Valent B, Khang CH. 2010. Recent advances in rice blast effector research. *Curr Opin Plant Biol* **13**: 434–441.
- van Damme M, Bozkurt TO, Cakir C, Schornack S, Sklenar J, Jones AM, Kamoun S. 2012. The Irish potato famine pathogen *Phytophthora infestans* translocates the CRN8 kinase into host plant cells. *PLoS Pathog* **8**: e1002875. doi: 10.1371/journal.ppat.1002875.
- Van den Ackerveken G, Marois E, Bonas U. 1996. Recognition of the bacterial avirulence protein AvrBs3 occurs inside the host plant cell. *Cell* **87**: 1307–1316.

- van den Burg HA, Harrison SJ, Joosten MH, Vervoort J, de Wit PJ. 2006. *Cladosporium fulvum* Avr4 protects fungal cell walls against hydrolysis by plant chitinases accumulating during infection. *Mol Plant Microbe Interact* **19**: 1420–1430.
- van der Hoorn RA, Kamoun S. 2008. From guard to decoy: A new model for perception of plant pathogen effectors. *Plant Cell* **20**: 2009–2017.
- van Esse HP, Van't Klooster JW, Bolton MD, Yadeta KA, van Baarlen P, Boeren S, Vervoort J, de Wit PJ, Thomma BP. 2008. The *Cladosporium fulvum* virulence protein Avr2 inhibits host proteases required for basal defense. *Plant Cell* **20**: 1948–1963.
- Wang Y, Li J, Hou S, Wang X, Li Y, Ren D, Chen S, Tang X, Zhou JM. 2010. A *Pseudomonas syringae* ADP-ribosyltransferase inhibits *Arabidopsis* mitogen-activated protein kinase kinases. *Plant Cell* **22**: 2033–2044.
- Wang Q, Han C, Ferreira AO, Yu X, Ye W, Tripathy S, Kale SD, Gu B, Sheng Y, Sui Y, et al. 2011. Transcriptional programming and functional interactions within the *Phytophthora sojae* RXLR effector repertoire. *Plant Cell* **23**: 2064–2086.
- Wei P, Wong WW, Park JS, Corcoran EE, Peisajovich SG, Onuffer JJ, Weiss A, Lim WA. 2012. Bacterial virulence proteins as tools to rewire kinase pathways in yeast and immune cells. *Nature* **488**: 384–388.
- Weiler EW, Kutchan TM, Gorba T, Brodschelm W, Niesel U, Bublitz F. 1994. The *Pseudomonas* phytotoxin coronatine mimics octadecanoid signalling molecules of higher plants. *FEBS Lett* **345**: 9–13.
- Westwood JH, Yoder JJ, Timko MP, dePamphilis CW. 2010. The evolution of parasitism in plants. *Trends Plant Sci* **15**: 227–235.
- Wu S, Lu D, Kabbage M, Wei HL, Swingle B, Records AR, Dickman M, He P, Shan L. 2011. Bacterial effector HopF2 suppresses *Arabidopsis* innate immunity at the plasma membrane. *Mol Plant Microbe Interact* **24**: 585–593.
- Xiang T, Zong N, Zou Y, Wu Y, Zhang J, Xing W, Li Y, Tang X, Zhu L, Chai J, et al. 2008. *Pseudomonas syringae* effector AvrPto blocks innate immunity by targeting receptor kinases. *Curr Biol* **18**: 74–80.
- Xie X, Yoneyama K. 2010. The strigolactone story. *Annu Rev Phytopathol* **48**: 93–117.
- Zeng L, Velasquez AC, Munkvold KR, Zhang J, Martin GB. 2012. A tomato LysM receptor-like kinase promotes immunity and its kinase activity is inhibited by AvrPtoB. *Plant J* **69**: 92–103.
- Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JD, Boller T, Felix G. 2006. Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-mediated transformation. *Cell* **125**: 749–760.

Cold Spring Harbor Symposia on Quantitative Biology

Effector Biology of Plant-Associated Organisms: Concepts and Perspectives

J. Win, A. Chaparro-Garcia, K. Belhaj, et al.

Cold Spring Harb Symp Quant Biol 2012 77: 235-247 originally published online December 6, 2012

Access the most recent version at doi:[10.1101/sqb.2012.77.015933](https://doi.org/10.1101/sqb.2012.77.015933)

References

This article cites 122 articles, 41 of which can be accessed free at:
<http://symposium.cshlp.org/content/77/235.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in
the box at the top right corner of the article or [click here](#).

To subscribe to *Cold Spring Harbor Symposia on Quantitative Biology* go to:
<http://symposium.cshlp.org/subscriptions>